

**The Synthesis and Biological Activity of N-Acylated Amino Acids.
A Collaborative Effort of Distributed Drug Discovery (D3)**

William L. Scott, Łukasz Popiołek,[†] Anna Biernasiuk,[‡]

Jake R. Hitchens, J. Geno Samaritoni, and Martin J. O'Donnell

[†] *Department of Organic Chemistry, Faculty of Pharmacy, Medical University of Lublin (Poland)*

[‡] *Department of Pharmaceutical Microbiology, Faculty of Pharmacy,
Medical University of Lublin (Poland)*

Indiana University Purdue University Indianapolis

As part of a Distributed Drug Discovery collaborative effort between students at IUPUI and Medical University of Lublin (Poland), the solid-phase combinatorial synthesis of a series of natural, acylated tyrosine (**1**) and phenylalanine (**2**) analogs was carried out in replicated fashion. The crude samples were purified and characterized by LC/MS, proton NMR, and in cases involving novel structures, by proton and carbon-13 NMR and high-resolution mass spectrometry. The samples were characterized in biological assays at the Medical University of Lublin against the Gram-positive bacteria *Staphylococcus aureus* ATCC 25923, *Staphylococcus epidermidis* ATCC 12228, *Bacillus cereus* ATCC 10876, *Bacillus subtilis* ATCC 10876, and *Micrococcus luteus* ATCC 10240. Although activity of the 2-nitro and 3-nitro derivatives of phenylalanine was not reproduced by the IUPUI samples, the 5-chlorosalicylic acid derivative **1g** demonstrated good activity against *M. luteus* (MIC = 62.5 µg/mL) and moderate activity against *S. aureus*, *S. epidermidis*, and *B. cereus*.

